

SOCIAL ANXIETY DISORDER: More Than Just Shyness



Are you extremely afraid of being judged by others?

Are you very self-conscious in everyday social situations?

Do you avoid meeting new people?

If you have been feeling this way for at least six months and these feelings make it hard for you to do everyday tasks—such as talking to people at work or school—you may have a **social anxiety disorder**.



National Institute
of Mental Health

Social anxiety disorder (also called social phobia) is a mental health condition. It is an intense, persistent fear of being watched and judged by others. This fear can affect work, school, and your other day-to-day activities. It can even make it hard to make and keep friends. But social anxiety disorder doesn't have to stop you from reaching your potential. Treatment can help you overcome your symptoms.

What is it like having social anxiety disorder?

In school, I was always afraid of being called on, even when I knew the answers. I didn't want people to think I was stupid or boring. My heart would pound and I would feel dizzy and sick. When I got a job, I hated to meet with my boss or talk in a meeting. I couldn't attend my best friend's wedding reception because I was afraid of having to meet new people. I tried to calm myself by drinking several glasses of wine before an event and then I started drinking every day to try to face what I had to do.

I finally talked to my doctor because I was tired of feeling this way and I was worried that I would lose my job. I now take medicine and meet with a counselor to talk about ways to cope with my fears. I refuse to use alcohol to escape my fears and I'm on my way to feeling better.

What is social anxiety disorder?

Social anxiety disorder is a common type of anxiety disorder. A person with social anxiety disorder feels symptoms of anxiety or fear in certain or all social situations, such as meeting new people, dating, being on a job interview, answering a question in class, or having to talk to a cashier in a store. Doing everyday things in front of people—such as eating or drinking in front of others or using a public restroom—also causes anxiety or fear. The person is afraid that he or she will be humiliated, judged, and rejected.

The fear that people with social anxiety disorder have in social situations is so strong that they feel it is beyond their ability to control. As a result, it gets in the way of going to work, attending school, or doing everyday things. People with social anxiety disorder may worry about these and other things for weeks before they happen. Sometimes, they end up staying away from places or events where they think they might have to do something that will embarrass them.

Some people with the disorder do not have anxiety in social situations but have performance anxiety instead. They feel physical symptoms of anxiety in situations such as giving a speech, playing a sports game, or dancing or playing a musical instrument on stage.

Social anxiety disorder usually starts during youth in people who are extremely shy. Social anxiety disorder is not uncommon; research suggests that about 7 percent of Americans are affected. Without treatment, social anxiety disorder can last for many years or a lifetime and prevent a person from reaching his or her full potential.

What are the signs and symptoms of social anxiety disorder?

When having to perform in front of or be around others, people with social anxiety disorder tend to:

- Blush, sweat, tremble, feel a rapid heart rate, or feel their “mind going blank”
- Feel nauseous or sick to their stomach
- Show a rigid body posture, make little eye contact, or speak with an overly soft voice

- Find it scary and difficult to be with other people, especially those they don't already know, and have a hard time talking to them even though they wish they could
- Be very self-conscious in front of other people and feel embarrassed and awkward
- Be very afraid that other people will judge them
- Stay away from places where there are other people

What causes social anxiety disorder?

Social anxiety disorder sometimes runs in families, but no one knows for sure why some family members have it while others don't. Researchers have found that several parts of the brain are involved in fear and anxiety. Some researchers think that misreading of others' behavior may play a role in causing or worsening social anxiety. For example, you may think that people are staring or frowning at you when they truly are not. Underdeveloped social skills are another possible contributor to social anxiety. For example, if you have underdeveloped social skills, you may feel discouraged after talking with people and may worry about doing it in the future. By learning more about fear and anxiety in the brain, scientists may be able to create better treatments. Researchers are also looking for ways in which stress and environmental factors may play a role.

How is social anxiety disorder treated?

First, talk to your doctor or health care professional about your symptoms. Your doctor should do an exam and ask you about your health history to make sure that an unrelated physical problem is not causing your symptoms. Your doctor may refer you to a mental health specialist, such as a psychiatrist, psychologist, clinical social worker, or counselor. The first step to effective treatment is to have a diagnosis made, usually by a mental health specialist.

Social anxiety disorder is generally treated with psychotherapy (sometimes called "talk" therapy), medication, or both. Speak with your doctor or health care provider about the best treatment for you. If your health care provider cannot provide a referral, visit the NIMH Help for Mental Illnesses web page at www.nimh.nih.gov/findhelp for resources you may find helpful.

Psychotherapy

A type of psychotherapy called cognitive behavioral therapy (CBT) is especially useful for treating social anxiety disorder. CBT teaches you different ways of thinking, behaving, and reacting to situations that help you feel less anxious and fearful. It can also help you learn and practice social skills. CBT delivered in a group format can be especially helpful. For more information on psychotherapy, please visit www.nimh.nih.gov/health/topics/psychotherapies.

Support Groups

Many people with social anxiety also find support groups helpful. In a group of people who all have social anxiety disorder, you can receive unbiased, honest feedback about how others in the group see you. This way, you can learn that your thoughts about judgment and rejection are not true or are distorted. You can also learn how others with social anxiety disorder approach and overcome the fear of social situations.

Medication

There are three types of medications used to help treat social anxiety disorder:

- Anti-anxiety medications
- Antidepressants
- Beta-blockers

Anti-anxiety medications are powerful and begin working right away to reduce anxious feelings; however, these medications are usually not taken for long periods of time. People can build up a tolerance if they are taken over a long period of time and may need higher and higher doses to get the same effect. Some people may even become dependent on them. To avoid these problems, doctors usually prescribe anti-anxiety medications for short periods, a practice that is especially helpful for older adults.

Antidepressants are mainly used to treat depression, but are also helpful for the symptoms of social anxiety disorder. In contrast to anti-anxiety medications, they may take several weeks to start working. Antidepressants may also cause side effects, such as headaches, nausea, or difficulty

sleeping. These side effects are usually not severe for most people, especially if the dose starts off low and is increased slowly over time. Talk to your doctor about any side effects that you have.

Beta-blockers are medicines that can help block some of the physical symptoms of anxiety on the body, such as an increased heart rate, sweating, or tremors. Beta-blockers are commonly the medications of choice for the “performance anxiety” type of social anxiety.

Your doctor will work with you to find the best medication, dose, and duration of treatment. Many people with social anxiety disorder obtain the best results with a combination of medication and CBT or other psychotherapies.

Don’t give up on treatment too quickly. Both psychotherapy and medication can take some time to work. A healthy lifestyle can also help combat anxiety. Make sure to get enough sleep and exercise, eat a healthy diet, and turn to family and friends who you trust for support.

For basic information about these and other mental health medications, visit **www.nimh.nih.gov/health/topics/mental-health-medications**.

Visit the Food and Drug Administration’s website (**www.fda.gov/**) for the latest information on warnings, patient medication guides, or newly approved medications.

For More Information

To learn more about social anxiety disorder, visit:

MedlinePlus (National Library of Medicine)

<http://medlineplus.gov>

(En Español: **<http://medlineplus.gov/spanish>**)

For information on clinical trials, visit:

ClinicalTrials.gov: <http://www.clinicaltrials.gov>

(En Español: **<http://salud.nih.gov/investigacion-clinica/>**)

For more information on conditions that affect mental health, resources, and research, visit the NIMH website **<http://www.nimh.nih.gov>**

Finding Help

Mental Health Treatment Program Locator

The Substance Abuse and Mental Health Services Administration (SAMHSA) provides this online resource for locating mental health treatment facilities and programs. The Mental Health Treatment Locator section of the Behavioral Health Treatment Services Locator lists facilities providing mental health services to persons with mental illness. Find a facility in your state at www.findtreatment.samhsa.gov/. For additional resources, visit www.nimh.nih.gov/findhelp.

Reprints

This publication is in the public domain and may be reproduced or copied without permission from NIMH. Citation of NIMH as a source is appreciated. We encourage you to reproduce it and use it in your efforts to improve public health. However, using government materials inappropriately can raise legal or ethical concerns, so we ask you to use these guidelines:

- NIMH does not endorse or recommend any commercial products, processes, or services, and our publications may not be used for advertising or endorsement purposes.
- NIMH does not provide specific medical advice or treatment recommendations or referrals; our materials may not be used in a manner that has the appearance of providing such information.
- NIMH requests that non-Federal organizations not alter our publications in ways that will jeopardize the integrity and “brand” when using the publication.
- The addition of non-Federal Government logos and website links may not have the appearance of NIMH endorsement of any specific commercial products or services, or medical treatments or services.
- Images used in publications are of models and are used for illustrative purposes only. Use of some images is restricted.

If you have questions regarding these guidelines and use of NIMH publications, please contact the NIMH Information Resource Center at **1-866-615-6464** or e-mail nimhinfo@nih.gov.

National Institute of Mental Health

Office of Science Policy, Planning, and Communications
Science Writing, Press, and Dissemination Branch

6001 Executive Boulevard

Room 6200, MSC 9663

Bethesda, MD 20892-9663

Phone: 301-443-4513 or 1-866-615-NIMH (6464) toll-free

TTY: 301-443-8431 or 1-866-415-8051 toll-free

Fax: 301-443-4279

Email: nimhinfo@nih.gov

Website: www.nimh.nih.gov



National Institute
of Mental Health

U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES

National Institutes of Health

NIH Publication No. QF 16-4678

Revised 2016

